

## Burnout Assessment

Formatted: Font: 12 pt

Most people have an intuitive idea of what burnout is. From the research perspective, burnout is one of the elements of compassion fatigue. It is associated with feelings of hopelessness and difficulties in dealing with work or in doing your job effectively. These negative feelings usually have a gradual onset. They can reflect the feeling that your efforts make no difference, or they can be associated with a very high workload or a non-supportive work environment. Higher scores on this scale mean that you are at higher risk for burnout. The average score on the burnout scale is 50 (SD 10; alpha scale reliability .75). About 25% of people score above 57 and about 25% of people score below 43. If your score is below 18, this probably reflects positive feelings about your ability to be effective in your work. If you score above 57 you may wish to think about what at work makes you feel like you are not effective in your position. Your score may reflect your mood; perhaps you were having a "bad day" or are in need of some time off. If the high score persists or if it is reflective of other worries, it may be a cause for concern.

## Compassion Fatigue

The second component of Compassion Fatigue (CF) is secondary traumatic stress (STS). It is about your work-related, secondary exposure to extremely or traumatically stressful events. Developing problems due to exposure to other's trauma is somewhat rare but does happen to many people who care for those who have experienced extremely or traumatically stressful events. For example, you may repeatedly hear stories about the traumatic things that happen to other people, commonly called Vicarious Traumatization. You may see or provide treatment to people who have experienced horrific events. If your work puts you directly in the path of danger, due to your work as a soldier or civilian working in military medicine personnel, this is not secondary exposure; your exposure is primary. However, if you are exposed to others' traumatic events as a result of your work, such as providing care to casualties or for those in a military medical rehabilitation facility, this is secondary exposure. The symptoms of STS are usually rapid in onset and associated with a particular event. They may include being afraid, having difficulty sleeping, having images of the upsetting event pop into your mind, or avoiding things that remind you of the event. The average score on this scale is 50 (SD 10; alpha scale reliability .81). About 25% of people score below 43 and about 25% of people score above 57. If your score is above 57, you may want to take some time to think about what at work may be frightening to you or if there is some other reason for the elevated score. While higher scores do not mean that you do have a problem, they are an indication that you may want to examine how you feel about your work and your work environment. You may wish to discuss this with your supervisor, a colleague, or a health care professional.

© B. Hudnall Stamm, 2009. Professional Quality of Life: Compassion Satisfaction and Fatigue Version 5 (ProQOL). /www.isu.edu/~bhstamm or www.proqol.org. This test may be freely copied as long as (a) author is credited, (b) no changes are made, and (c) it is not sold.